

PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM	Books
Search PubMed	<input type="checkbox"/> for					Go	Clear	
Limits Preview/Index History Clipboard Details								

Entrez
PubMed

☐ 1: Biochemistry 1993 Jun 15;32(23):6116-20

[Related Articles](#), [NEW Books](#)

Rapid kinetic analysis of a point mutant of HIV-1 reverse transcriptase lacking ribonuclease H activity.

Dudding LR, Mizrahi V.

South African Institute for Medical Research, Department of Haematology, University of the Witwatersrand Medical School, Johannesburg.


The comparative kinetics of RNA-dependent DNA polymerization catalyzed by wild-type HIV-1 reverse transcriptase and a point mutant specifically lacking RNase H activity were analyzed using a heteropolymeric substrate consisting of a 19-mer primer hybridized to a 345-nucleotide RNA template. The rapid-quench product distributions generated under single-turnover conditions, in which primer extension by the two enzymes was restricted to the incorporation of 5 nucleotides (N+5), were significantly different. Whereas the wild-type enzyme catalyzed synthesis of the N+5 product over the time course of the reaction (20 ms-10 s) with a relatively low degree of processivity, the extent of accumulation of the intermediate N+2 and N+3 products was grossly exaggerated in the parallel mutant-catalyzed time course. The observation of concomitant polymerase-dependent hydrolysis during the course of synthesis catalyzed by the wild-type enzyme suggested that the inability of the RNase H- mutant to hydrolyze the RNA template created blocks to further synthesis by reducing the rates of DNA polymerization at these intermediate positions, and hence impaired the ability of this mutant to complete cDNA synthesis.

PMID: 7685188 [PubMed - indexed for MEDLINE]

Display	Abstract	<input type="checkbox"/>	Sort	<input type="checkbox"/>	Save	Text	Clip Add	Order
---------	----------	--------------------------	------	--------------------------	------	------	----------	-------

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Freedom of Information Act](#) | [Disclaimer](#)

spare-sun-solaris2.8 Dec 20 2001 12:01:15



PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM	Books
Search	PubMed	<input type="checkbox"/> for					Go	Clear
		Limits	Preview/Index	History	Clipboard	Details		

Entrez
PubMed

Display	Abstract	<input type="checkbox"/>	Sort	<input type="checkbox"/>	Save	Text	Clip Add	Order
---------	----------	--------------------------	------	--------------------------	------	------	----------	-------

☐ 1: Mol Biotechnol 1997 Aug;8(1):61-77Related Articles, ^{NEW} BooksPubMed
Services

Reverse transcriptase. The use of cloned Moloney murine leukemia virus reverse transcriptase to synthesize DNA from RNA.

Gerard GF, Fox DK, Nathan M, D'Alessio JM.

Life Technologies, Rockville, MD 20850, USA. ggerard@lifetech.com

Related
Resources


Reverse transcriptase (RT) is the key enzyme required for conversion of RNA to DNA. Cloning of Moloney murine leukemia virus (MMLV) RT has enable engineering an RT that lacks endogenous RNase H activity. RT catalyzes cDNA synthesis more efficiently in the absence of RNase H. We describe here a number of properties of MMLV RT and RNase H-minus MMLV RT not summarized in a single location elsewhere, providing a basis for best use of these enzymes in cDNA synthesis. In addition, general guidelines and detailed protocols are provided for use of MMLV RTs in one tube double-stranded cDNA synthesis, in [32P]cDNA synthesis, and in RT-PCR and long RT-PCR.

PMID: 9327398 [PubMed - indexed for MEDLINE]

Display	Abstract	<input type="checkbox"/>	Sort	<input type="checkbox"/>	Save	Text	Clip Add	Order
---------	----------	--------------------------	------	--------------------------	------	------	----------	-------

Write to the Help Desk
NCBI | NLM | NIH
Department of Health & Human Services
Freedom of Information Act | Disclaimer

spare-sun-solaris2.8 [Dec 20 2001 12:01:15]



PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM	Books
Search PubMed	▼ for						Go	Clear
Limits		Preview/Index		History		Clipboard		Details

Entrez
PubMed

Display	Abstract	▼	Sort	▼	Save	Text	Clip Add	Order
---------	----------	---	------	---	------	------	----------	-------

☐ 1: Biotechniques 1997 Jun;22(6):1119-22, 1124-6Related Articles, ^{NEW} Books

Analysis of the size distribution of first-strand cDNA molecules.

PubMed
Services**Bodescot M, Brison O.**

Institut Gustave-Roussy Villejuif, France.

Related
Resources


We have developed a method to analyze the size distribution of the first-strand cDNA molecules corresponding to given mRNA species. First-strand molecules synthesized from cytoplasmic polyadenylated RNAs are separated by electrophoresis on an alkaline agarose gel, and a Southern blot hybridization is performed. As an example, we analyzed the first-strand molecules corresponding to the human c-myc mRNAs. This method can be used to determine whether full-length, first-strand molecules corresponding to an mRNA species to be cloned are synthesized efficiently. Interestingly, this method allows one to analyze full-length, first-strand cDNA molecules with a much higher resolution than Northern blot analysis of mRNA molecules. This method can therefore be used to discriminate between the multiple mRNA species transcribed from a given gene or the homologous mRNA species transcribed from a given gene family.

PMID: 9187762 [PubMed - indexed for MEDLINE]

Display	Abstract	▼	Sort	▼	Save	Text	Clip Add	Order
---------	----------	---	------	---	------	------	----------	-------

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Freedom of Information Act](#) | [Disclaimer](#)

spare-sun-solaris2.8 Dec 20 2001 12:01:15



PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM	Books	
Search	PubMed	▼	for					Go	Clear
		Limits	Preview/Index	History	Clipboard	Details			
Display		Abstract	▼	Sort	▼	Save	Text	Clip Add	Order

Entrez
PubMed

☐ 1: Biochemistry 1983 May 10;22(10):2365-72

Related Articles, NEW Books

PubMed
Services


Reverse transcriptase and its associated ribonuclease H: interplay of two enzyme activities controls the yield of single-stranded complementary deoxyribonucleic acid.

Berger SL, Wallace DM, Puskas RS, Eschenfeldt WH.

Related
Resources

The synthesis of single-stranded globin cDNA by the RNA-directed DNA polymerase activity of reverse transcriptase in the presence of oligothymidylate primers was investigated in order to determine the limitations to higher yields. The results indicated that the associated ribonuclease H activity, an integral part of reverse transcriptase, plays a large role in the synthesis of the first strand of cDNA and that the interplay of the two enzyme activities for any specific set of conditions determines the yield of single-stranded products. In both the presence and the absence of polymerization, the associated ribonuclease H catalyzed the deadenylation of mRNA, producing molecules that were somewhat shorter, highly homogeneous in size, and fully translatable into globin protein. They were also entirely lacking in the ability to serve as templates for cDNA synthesis. The reaction was completely dependent on oligothymidylate and completely independent of deoxyribonucleoside triphosphates. The initial rate of deadenylation was one-fourth the initial rate of initiation of polymerization when saturating levels of deoxyribonucleoside triphosphates were used in the polymerase reaction. In the presence of ribonuclease H activity, the DNA polymerase catalyzed the synthesis of an array of cDNAs including some that were full length. The initiation of polymerization was rate limiting: once synthesis had begun, it required 1-1.5 min to transcribe globin mRNA. However, most primers that were elongated were aborted prematurely. Maximum synthesis of full-length cDNA required stoichiometric levels of enzyme and high triphosphate levels, but regardless of conditions, the sum of completed cDNA and deadenylated mRNA accounted for only 50% of the input mRNA. The data fit a model in which synthesis of full-length cDNA molecules depends on the arrangement of primers and transcription initiation complexes on the poly(A) "tail" of mRNA.

PMID: 6190507 [PubMed - indexed for MEDLINE]



PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM	Books
Search PubMed	<input type="checkbox"/> for						Go	Clear
Limits		Preview/Index		History		Clipboard		Details

Entrez
PubMed

☐ 1: Proc R Soc Lond B Biol Sci 1991 Mar
22;243(1308):235-9

Related Articles, ^{NEW} Books

PubMed
Services

Reverse-transcriptase-associated RNaseH activity mediates template switching during reverse transcription in vitro.

Garces J, Wittek R.

Institut de Biologie Animale, Universite de Lausanne, Switzerland.

Related
Resources


During the first steps of reverse transcription of the retroviral genome, sequences present at the extremities of the RNA are used to reconstitute a host cell PolIII promoter. The assembly of the promoter occurs by template switching, which takes advantage of a direct repeat at the ends of the RNA molecule. These steps are catalysed by the viral reverse transcriptase, which carries an intrinsic RNaseH activity that is probably also involved therein. To study the role of the RNaseH activity in this first template-switching event, an in vitro system has been developed based on primer extensions of synthetic RNAs. When an RNA was reverse transcribed with wild-type reverse transcriptase in the presence of a second RNA the 3' part of which was repeated at the 5' end of the first one, extension products could be observed corresponding to a chimeric cDNA comprising both RNA species. This template switching could not be detected when a mutant reverse transcriptase lacking the RNaseH activity was used. The results show that the RNaseH activity is needed to remove the 5' RNA sequences from the cDNA:RNA hybrid thereby enabling its translocation to another RNA containing an appropriate complementary target sequence.

PMID: 1711233 [PubMed - indexed for MEDLINE]

Display	Abstract	<input type="checkbox"/>	Sort	<input type="checkbox"/>	Save	Text	Clip Add	Order
---------	----------	--------------------------	------	--------------------------	------	------	----------	-------

Write to the Help Desk
NCBI | NLM | NIH
Department of Health & Human Services
Freedom of Information Act | Disclaimer

spare-sun-solaris2.8 Dec 20 2001 12:01:15



PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM	Books
Search PubMed	▼ for						Go	Clear
Limits		Preview/Index		History	Clipboard	Details		

Entrez
PubMed

Display	Abstract	▼	Sort	▼	Save	Text	Clip Add	Order
---------	----------	---	------	---	------	------	----------	-------

☐ 1: J Biol Chem 1995 Oct 13;270(41):24135-45 Related Articles, NEW Books, LinkOut

FREE full text article at
www.jbc.org

PubMed
Services

Cleavage specificities of Moloney murine leukemia virus RNase H implicated in the second strand transfer during reverse transcription.

Schultz SJ, Whiting SH, Champoux JJ.

Department of Microbiology, School of Medicine, University of Washington, Seattle 98195-7242, USA.

Related
Resources

Reverse transcription of a retroviral RNA genome requires two template jumps to generate the linear double-stranded DNA required for integration. The RNase H activity of reverse transcriptase has several roles during this process. We have examined RNase H cleavages that define the maximal 3' and 5' ends of Moloney murine leukemia virus minus strand DNA prior to the second template jump. In both the endogenous reaction and on model substrates in vitro, RNase H cleaves the genomic RNA template between the second and third ribonucleotides 5' of the U5/PBS junction, but other minor cleavages between 1 and 10 nucleotides 5' of this junction are also observed. Similar experiments examining the specificity of RNase H for tRNA primer removal revealed that cleavage generally leaves a ribo A residue at the 5' end of minus strand DNA. These observations suggest that three bases are typically duplicated on the ends of the minus strands, leading to an intermediate following the second jump which contains unpaired nucleotides. Model substrates mimicking the structure of this intermediate demonstrate that reverse transcriptase has little difficulty in utilizing such a branched structure for the initiation of displacement synthesis.

PMID: 7592616 [PubMed - indexed for MEDLINE]

Display	Abstract	▼	Sort	▼	Save	Text	Clip Add	Order
---------	----------	---	------	---	------	------	----------	-------

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Freedom of Information Act](#) | [Disclaimer](#)